TRANSCRANIAL ECOCOLOR DOPPLER
from methodology to clinical applications

Transcranial Ecocolor Doppler is currently the elective Imaging method for noninvasive study of intracranial arteries and their hemodynamics. Feasible in adults and children, it is indicated for the early diagnosis of acute brain disease and in the follow-up of chronic cerebrovascular states.

TCD: EXAMINATION TECHNIQUE

With the patient in the supine position and with the operator behind the patient's head, it is possible to visualize the intracranial vessels through the 3 *acoustic windows*, areas of the skull, made up of the bones that allow the penetration of the US:

Ultrasound windows: Temporal, Occipital, Frontotemporal

Among them, the most interesting for diagnostic use is the temporal window (Fig. 1,2).

Fig. 1


Fig. 2

Rectangular area on the squamous part of the temporal bone subdivided in 3 zones: front, middle and rear.

Middle cerebral artery (MCA)

It collects about 60-70% of the blood coming from the *internal carotid arteries (ICA)*, allowing, therefore, the estimate of blood flow to the hemisphere ipsilateral (Fig. 3).
It allows to explore (Fig. 4):


Fig. 3


Fig. 4

Middle cerebral arteries (MCA);
in the M1 and M2 *Anterior cerebral arteries (ACA)* sections;
in A1 and A2, *Posterior cerebral arteries (PCA)*, P1 and P2.

In most patients, identification is possible through the 'butterfly wings sign' (Fig. 5).
In intracranial vessels, the Doppler signal highlights a flow with a predominant diastolic component (Fig. 6).

Fig. 5


Fig. 6

TCD: INDICATIONS

From the spectral analysis of the Doppler signal it is possible to calculate the cerebral flow rate.
Cerebral (Mean CBFV) (Table 1, 2)

Table 1 - Factors influencing cerebral blood flow velocity

Factor	Change in CBFV
Age	Increase up 6-10 yr then decrease
Sex	Women > men
Pregnancy	Decrement in the III Trimester
Hematocrit	Increase with decreasing Hct
PCO_2	Increase with increasing PCO_2
Pain	Arterial pressure increase with increasing MAP

CBFV: Cerebral blood flow velocity; MAP: Mean arterial pressure.

Table 2 - Mean cerebral blood flow (cm/s) related to age

Artery	Age 20-40 yr	Age 40-60 yr	Age > 60 yr
Anterior cerebral artery	56-60	53-61	44-51
Middle cerebral artery	74-81	72-73	58-59
Posterior cerebral artery P1	48-57	41-56	37-47
Posterior cerebral artery P2	43-51	40-57	37-47
Vertebral artery	37-51	29-50	30-37
Basilar artery	39-58	27-56	29-47

Mean CBFV = (PSV + [EDV × 2])/3

PSV = peak systolic velocity
EDV = end-diastolic blood flow velocity
Change of CBFV mean, may suggest vasospams

Lindegaard Ratio (LR): (Table 3)

Table 3 - Intracranial arteries: severity of vasospasm

	MFV	LR modified
MCA or ICA vasospasm (%)		
Mild (< 25)	120-140	3-6
Moderate (25-50)	15-199	3-6
Severe (> 50)	> 200	> 6
BA vasospasm (%)		
Possible vasospasm	70-85	2-2.49
Moderate (25-50)	> 35	2.5-2.99
Severe (> 50)	> 85	> 3

MCA: Middle cerebral artery; ICA: Internal carotid artery; LR: Lindegaard ratio; BA: Basilar artery; MFV: Mean flow velocity; MCA: Middle cerebral artery; ICA: Internal carotid artery; LR: Lindegaard ratio; BA: Basilar artery; MFV: Mean flow velocity.

MCA mean CBFV/extracranial ICA mean CBFV (v.n. <3)

BA mean CBFV/left or right extracranial VA mean CBFV (v.n. <2)

Fig. 7

TCD: CRYPTOGENIC STROKE AND t-RLS

In subjects aged < 55 years, in 40% of cases stroke is of cryptogenic origin. In these pts, the prevalence of Patent Foramen Ovale (PFO), responsible for paradoxical embolism due to *transient Right to Left shunt (t-RLS)*, is about 40%.

Examination technique (Fig. 7)
t-RLS is highlighted through shaken saline solution that behaves as a contrast.
Contrast agent: 9 mL of normal saline solution with 1 mL of air or blood, shaken up about 10 times through a system constituted by two 10 mL syringes linked by a 3-way stopcock. The agitated solution is administrated into the anticubital vein by an 18-gauge. The patient is then invited to perform a forced expiration against the closed glottis for a minimum of 10 seconds.

Table 4 - Grade of transient right to left shunting based on microembolic signals grading score

Grade transient shunt	MES
No shunt	0
Low grade shunt	1-10
Moderate grade shunt	11-25
High grade shunt	> 25 (showers) or uncountable (curtain effect)

MES: Microembolic signals

C-TCD, placing the PW Doppler slider on the MCA, show the passage of 'Microembolic Signals-MES', indirect manifestations of t-RLS.



Contrast-enhanced transcranial Doppler as a first-line screening test for the evaluation of a cryptogenic ischemic stroke. TCD: Transcranial Doppler; eTCD: Contrast-enhanced TCD; TEE: Transesophageal echocardiography; PFO: Patent foramen ovale; ASA: Acetylsalicylic acid; VKA: Vitamin K antagonist; MES: Microembolic signals; RRR: Recurrent risk reduction; CI: Contraindication.

Transcranial Doppler Ultrasound: Physical Principles and Principal Applications. Ballotta E, Da Giau G, Renon D, Furlan A, Meneghetti G, Baracchini C. J Cardiovasc Technol. 2004 Apr-Jun; 23(2):120-41. doi: 10.1177/089431800402300204. PMID: 15141184. Review. No abstract available.

Transcranial Doppler Ultrasound: Incremental Diagnostic Role in Cryptogenic Stroke. Ballotta E, D'Andrea A, Cecconi M, Angelini L, Bonetti R, Cavallini E, Pianella G, Casellato M, Da Giau G, Baracchini C, Meneghetti G, Thiene G, Basso C. J Cardiovasc Technol. 2014 Jul-Aug; 33(3):171-77. doi: 10.1016/j.jocs.2013.11.005. Epub 2013 Dec 18. Review.

Antonello D' Andrea, Marianna Conte, Marco Magliano

esaote